

# WHEN TO RETURN **IMMEDIATELY**



#### **BRING ANY SICK CHILD**







If not able to drink

If becomes sicker 
If develops a fever

#### BRING CHILD with DIARRHOEA





#### **BRING CHILD with COUGH**





If difficult breathing If fast breathing

### BRING YOUNG INFANT (<2 months)

Breastfeeding or drinking poorly Becomes sicker Develops a fever or feels cold to touch Fast breathing Difficult breathing

Yellow palms and soles (if infant has jaundice)

Diarrhoea with blood in stool

Name:	M F Date of Birth:
Address:	
-	Always bring this card with you to the clinic.

## **FLUIDS** FOR ANY SICK CHILD

- · Breastfeed frequently.
- Increase fluid. Give soup, rice water, yoghurt drinks, or clean water.

#### FOR CHILD WITH DIARRHOEA

Giving more fluid can be lifesaving! • Give these extra fluids, as

- much as the child will take:
  - ORS Solution
  - Food based fluids, such as: soup,

rice water, youghurt drinks

- Clean water
- Breastfeed more frequently and longer at each feeding.
- Continue giving extra fluids until diarrhoea stops.

BCG	DPT 1	DPT 2	DPT 3	DPT (Booster)	DT
OPV 0	OPV 1	OPV 2	OPV 3	OPV	IFA
OF V 0		OF V 2	OF V 3	OF V	
HE	 P B-1				
	HEP E	3 -2 HE	P B -3	MEASLES	VITAMIN A
urn for nevt im	munization or vita	amin A or IFA	supplementati	on on:	